


# A measurement of the parity-violating gamma-ray asymmetry in the radiative $n$ - $p$ capture




Seppo Penttila  
LANL

For the NPDGamma Collaboration

CGS 11  
Pruhonice  
September 2-6, 2002



The NPDGamma studies weak interaction between neutrons and protons in the  $\vec{n} + p \rightarrow d + \gamma$  (2.2 MeV) reaction. The experiment will measure  $A_\gamma$ , the parity-violating asymmetry in the distribution of emitted  $\gamma$ 's



$PV \rightarrow$  signature of the weak interaction, in the strong interaction parity is conserved.

Expected asymmetry  $A_\gamma \approx -5 \times 10^{-8}$

Goal experimental error  $\leq 0.5 \times 10^{-8}$

$A_\gamma$  is a clean measurement of  $H_\pi^1$  since

2-body system - no nuclear structure uncertainties

$$A_\gamma \approx -0.045 \quad H_\pi^1$$

## Weak Interaction

---

- Standard model specifies well how point like leptons and quarks interacting weakly by exchanging a weak boson;  $W^+$ ,  $W^-$ ,  $Z^0$ , ....
- We do not have a good picture of the weak interaction between hadrons.
- The strong interaction binds nucleons together to form nuclei, and is thus the primary interaction between protons and neutrons. Nucleon interactions take place on a scale of  $1/m_\pi \approx 1.5$  fm, short range repulsion.
- The masses of the weak bosons are large and therefore their Compton wavelengths are in range of  $1/M_W \approx 10^{-3}$  fm which is very small compared to the distances characterizing low-energy  $N$ - $N$  interactions.

At low energies weak interaction between nucleons cannot be explained by a simple  $Z$  or  $W$  - exchange. The picture is complicated by strong interaction.

The ratio of weak and strong amplitudes is  $4\pi G_F m_\pi^2 / g_{\pi NN}^2 \approx 10^{-7}$

## Low-Energy Hadronic Weak Interaction: $\Delta I=1$ neutral current component

---

- At low energies hadronic weak interaction can be considered as a point interaction of two currents; the charged and neutral weak currents

$$H_W = \frac{G_F}{\sqrt{2}} (J_W^+ J_W + J_Z^+ J_Z) = H_L + H_{SL} + H_{NL} (\Delta S = 1) + H_{NL} (\Delta S = 0)$$

- From isospin structure of the weak interaction for strangeness conserving  $\Delta S = 0$  part is obtained:
  - Charged current weak  $N$ - $N$  interaction components are  $\Delta I = 0$  and 2 but not to  $\Delta I = 1$ .
  - Neutral currents components are  $\Delta I = 0, 1, 2$ .

$\Rightarrow$  The neutral current should dominate the isospin  $\Delta I = 1$  channel.

$\Rightarrow A_\gamma$  in  $\vec{n} + p \rightarrow d + \gamma$  is a measure of the  $\Delta S = 0, \Delta I = 1$  part of the hadronic weak interaction and thus neutral current part of the interaction.

The main goals of the experiments are:

- to isolate the neutral current and
- to understand the mechanism by which the weak force is communicated over the long distances of  $N$ - $N$  interactions.

## The Low-Energy $N$ - $N$ Weak Interaction: one-meson exchange model

The low-energy  $N$ - $N$  weak interaction is conventionally described in a one - meson - exchange model, where one meson - nucleon vertex is weak and the other strong. This long - distance mechanism dominates at nuclear densities.

The six weak  $PV$  couplings;

$$H_{\pi}^1, H_{\rho}^0, H_{\rho}^1, H_{\rho}^{1'}, H_{\rho}^2, H_{\omega}^0, H_{\omega}^1$$

characterize the strengths of


isovector  $\pi$

isoscalar/isovector/isotensor  $\rho$

isoscalar/isovector  $\omega$


weak meson-nucleon couplings.

Measured in various combinations by a variety of observables.



$A_{\gamma} \approx -0.045 H_{\pi}^1$ . The  $\pi^{\pm}$  carries the important long-range part of the hadronic weak interaction, just as it does for the strong interaction.

# Simple Level Diagram of $n$ - $p$ System; $\bar{n} + p \rightarrow d + \gamma$ is primarily sensitive to the $\Delta I = 1$ component of the weak interaction



Mixing amplitudes:

$$\langle ^3S_1 | V_W | ^3P_1 \rangle; \Delta I = 1$$

$$\langle ^3S_1 | V_W | ^1P_1 \rangle; \Delta I = 0$$

$$\langle ^1S_0 | V_W | ^3P_0 \rangle; \Delta I = 2$$

- Weak interaction mixes in  $P$  waves to the singlet and triplet  $S$ -waves in initial and final states.
- Parity conserving transition is  $M1$ .
- Parity violation arises from mixing in  $P$  states and interference of the  $E1$  transitions.
- $A_\gamma$  is coming from  $^3S_1 - ^3P_1$  mixing and interference of  $E1$ - $M1$  transitions -  $\Delta I = 1$  channel.

Measurement of the Parity - Violating Gamma Asymmetry  $A_\gamma$  in the  
Capture of Polarized Cold Neutrons by Para-Hydrogen,  $\bar{n} + p \rightarrow d + \gamma$

**NPDGamma Collaboration**

J. D. Bowman (Spokesperson), G. S. Mitchell,  
S. I. Penttila, W. S. Wilburn, and V. W. Yuan

Los Alamos National Laboratory

G. L. Jones

Hamilton College

M. Gericke, H. Nann, and W. M. Snow

Indiana University

E. I. Sharapov

Joint Institute for Nuclear Research,  
Dubna, Russia

S. Ishimoto, Y. Masuda, and K. Morimoto

KEK National Laboratory, Japan

T. R. Gentile

National Institute of Standards and  
Technology

R. D. Carlini

Thomas Jefferson National Accelerator  
Facility

T. Case, S. J. Freedman

University of California, Berkeley

T. B. Smith

University of Dayton

S. A. Page and W. D. Ramsay

University of Manitoba and TRIUMF,  
Canada

T. E. Chupp and K. P. Coulter

University of Michigan

B. Hersman and M. B. Leuschner

University of New Hampshire


G. L. Greene

University of Tennessee

## The $\vec{n} + p \rightarrow d + \gamma$ Experiment is Designed for the LANSCE Pulsed Cold Spallation Neutron Source

The goal experimental error of  $0.5 \times 10^{-8}$  is a challenge for a polarized cold neutron flux from a spallation source as well as for the control of systematic errors.


- Some  $4 \times 10^{17}$  neutrons are required for statistics.
- Neutron time-of-flight used to design the experiment with systematic errors less than  $0.5 \times 10^{-8}$ .



The flight path 12 at LANSCE has a peak flux of  $2 \times 10^7$  n/cm<sup>2</sup>/s at about 8 meV. Neutron pulse rate is 20 Hz  $\rightarrow$  *TOF* frame = 50 ms. To reach the statistics of  $0.5 \times 10^{-8}$  eight months of running is needed.



# NPDGamma Experimental Setup



$$\frac{d\omega}{d\Omega} = \frac{1}{4\pi} (1 + A_\gamma \cos(\Theta \sigma_n \cdot k_n))$$

## Neutrons Polarized by Optically-Polarized $^3\text{He}$ Spin Filter


$$P_n = \tanh[nl\sigma_p(E_n)P_{\text{He}}]$$

$$T_n = T_n^0 \cosh[nl\sigma_p(E_n)P_{\text{He}}]$$


$$P_n = (1 - (T_n^0/T_n^p)^2)^{1/2}$$

$^3\text{He}$  neutron spin filter:

- In a  $^3\text{He}$  cell Rb atoms are polarized by laser light. Through spin exchange  $^3\text{He}$  gas is nuclear polarized.
- Cross section of the  $n$ - $^3\text{He}$  singlet state is much larger than the triplet state.
- Therefore, neutrons with spin antiparallel with  $^3\text{He}$  spins are absorbed and neutrons with spin parallel with  $^3\text{He}$  spins are transmitted  $\rightarrow$  neutron spin filter



## $^3\text{He}$ Spin-Filter Setup



- Large area, 11 cm in dia,  $^3\text{He}$  cells are required to cover the beam.
- $^3\text{He}$  spin filter allows a compact experimental setup.
- $^3\text{He}$  spin filter offers an extra spin flip without a change of the  $B$  field.

A 11-cm in diameter cell has  $T_1 > 500$  hr.  $^3\text{He}$  polarization 52% has been measured.



100 W light at 795 nm


## RF Neutron Spin Flipper: spin reversal in broad neutron energy range

- Neutron spin is guided by a  $B_0 = 10$  gauss field.
- Magnetic gradients have to be  $< 1$  mgauss/cm - no Stern-Gerlach steering  $\rightarrow$  false asymmetry
- RF spin flipper (RFSF) is the main control of systematic errors. Spin flip at 20 Hz.
- Magnetic field fluctuations less than 20 mgauss.
- Spin reversal with a RF field.
  - $E_n$  is proportional to  $1/(tof)^2$
  - At resonance a tilt angle of the spin is  $\Theta = \gamma B_1 \Delta t$
  - To precess a neutron spin by  $\Theta = \pi$ 

$$B_1 = (\pi L \gamma d) / (tof)$$
  - Spin-flip efficiency  $> 95\%$  achieved.

Other spin flips are guide field or  $^3\text{He}$  polarization


## RF Spin Flipper; spin-flip efficiency





## 20-liter Liquid Para-Hydrogen Target


- To maintain neutron spin in scattering a para- hydrogen target is required.
- The 30 cm in diameter and 30 cm long target captures 60% of incident neutrons.
- At 17 K only 0.05% of LH<sub>2</sub> is in ortho state → 1% of incident neutrons will be depolarized.
- Target cryostat materials selected so that false asymmetries  $< 10^{-10}$ .



Neutron mean free paths at 4 meV in


- ortho-hydrogen is  $\lambda \approx 2$  cm,
- para-hydrogen is  $\lambda \approx 20$  cm
- for a  $n$ - $p$  capture is  $\lambda \approx 50$ cm.

## A safe 20-liter Liquid Para-Hydrogen Target



## CsI(Tl) Gamma Detector


- Up-down  $\gamma$  – asymmetry will be measured.
- The detector has to be aligned with the  $B_0$  guide field better than 20 mrad.
- CsI(Tl) was selected because of :
  - large number of photoelectrons, measured  $>1000/\text{MeV } \gamma\text{-ray}$
  - interaction length of a 2.2-MeV  $\gamma\text{-ray}$  in CsI is about 5 cm  $\rightarrow$  95% of  $\gamma$ 's will be stopped in 15 cm.
- The gamma detector has 48 CsI modules -  $15 \times 15 \times 15 \text{ cm}^3$ . Total of 0.7 metric ton of CsI(Tl).



48 CsI(Tl) detectors have been received and are under testing.



## CsI(Tl) Gamma Detector




Detector solid angle is  $\approx 3\pi$ .

## Current Mode Detection with Vacuum Photo Diodes


- Rates are about  $10^{10} \gamma/\text{s} \rightarrow$  current mode detection.
- Light from CsI(Tl) is detected by 3" vacuum photodiodes which have S20 photocathodes.
  - vacuum photodiode linearity  $< 10^{-4}$  and
  - magnetic field sensitivity  $< 10^{-4}/\text{G}$  and  $< 10^{-5}/\text{G}^2$
- Vacuum photodiode has a gain of 1  $\rightarrow$  a high-gain low-noise  $I \rightarrow V$  preamplifier is required.
- In the current mode detection statistical fluctuations (counting statistics) appear as a shot noise in photodiode current.
- RMS of the shot noise current density is  $i_{\text{noise}}/\sqrt{f} = \sqrt{2qI} \approx n_{\text{pe}}e\sqrt{2 \times \text{rate}} \approx 5 \text{ pA}/\sqrt{\text{Hz}}$  per detector.

## Counting statistics vs electrical noise




- Sources of noise; Johnson noise in  $R_f$ , current and voltage noises in the op amp input, and photocathode dark current
- If  $R_f = 60 \text{ M}\Omega$  then the total calculated equivalent noise at the input is  $19 \text{ fA}/\sqrt{\text{Hz}}$  dominated by  $R_f$ .
- Noise value measured from the circuit is about  $20 \text{ fA}/\sqrt{\text{Hz}}$ .  
→ counting statistics/noise  $\approx 250$ .
- Counting statistics rules the run time.
- To measure beam-off false asymmetry to  $0.5 \times 10^{-8}$  beam-off time of a few minutes is needed.

# $n+p \rightarrow d+\gamma$ Experiment Layout



# Systematic Errors

- Physics - correlated with neutron spin:
  - activated materials - emit  $\gamma$ s in  $\beta$ -decay
  - Stern-Gerlach steering
  - L-R asymmetry
    - $n - p$  elastic scattering
    - $n - p$  parity allowed asymmetry
    - Mott-Schwinger scattering
- Instrumental sources
  - electronics, stray magnetic fields, gain stability
- Monitoring:
  - Null test at  $E_n > 15$  meV and at end of each pulse.



## Summary

- The  $\vec{n} + p \rightarrow d + \gamma$  reaction is the most sensitive way to isolate the neutral current component in the hadronic weak interaction and to determine  $H_{\pi}^1$ .
- With pulsed cold neutrons  $A_{\gamma}$  can be measured with the statistical precision of  $5 \times 10^{-8}$ 
  - *In situ* control of systematic errors by *TOF* information.
- The  $\vec{n} + p \rightarrow d + \gamma$  experiment will use  $^3\text{He}$  neutron spin filter, RF spin flipper, and current mode detection.
- Commissioning of the experiment will start in spring 2003.
- An interesting follow-up experiment could be  $\vec{n} + d \rightarrow t + \gamma$ .

